

Supplementary material to the paper Wells AF et al. Vegetation classification for northwestern Arctic Alaska using an EcoVeg approach: tussock tundra and low and tall willow groups and alliances. Vegetation Classification and Survey. DOI: 10.3897/VCS.65469

Supplementary material 4. Key to the low/tall willow and tussock tundra plant associations from the Ecological Land Survey Legacy Database (ELD) Arctic Plant Association Classification, Alaska.

1a. The site is located in the Arctic or Boreal ecoregion and is not influenced by salt water. The soils are flooded or wet for most or all of the freeze-free period. The vegetation is characterized by hydrophytic plants with at least 10% cover of vascular plants. Arctic & Boreal Freshwater Marsh, Wet Meadow & Shrubland Division (2.C.4.Np).....	2
2a. Site is located in the Boreal ecoregion. Not considered further here.....	
..... North American Boreal Marsh, Wet Meadow & Shrubland Macrogroup (M894).	
2b. Site is located in the Arctic ecoregion. Arctic Freshwater Marsh, Wet Meadow & Shrubland Macrogroup (M870).	3
3a. Combined cover of low (0.2-1.5 m) and tall (≥ 1.5 m) shrubs $\geq 25\%$ or combined cover of low, tall, and true dwarf shrubs $\geq 35\%$. North American Arctic Wet Shrubland Group (G830).	4
4a. Sites located in ombrotrophic wetlands, i.e., hydrology controlled predominantly by precipitation.....	5
5a. Combined cover of low shrubs $\geq 25\%$. Arctic Ombrotrophic Wet Low Shrublands Alliance Proposed (A4366p)..	6
6a. <i>Salix pulchra</i> is the dominant low shrub.....	7
7a. <i>Betula nana</i> cover $\geq 10\%$	8
8a. Combined cover of <i>Carex aquatilis</i> and <i>Eriophorum angustifolium</i> $\geq 10\%$	
..... <i>Betula nana</i> - <i>Salix pulchra</i> / <i>Carex aquatilis</i> - <i>Eriophorum angustifolium</i>	
8b. Vegetation not as above.....	Undefined Association
7b. <i>Betula nana</i> cover $< 10\%$	9
9a. <i>Carex aquatilis</i> and <i>Eriophorum angustifolium</i> are both present at $\geq 3\%$ each, and the combined cover of <i>C. aquatilis</i> and <i>E. angustifolium</i> is $\geq 10\%$	10
10a. <i>Sphagnum</i> cover $\geq 10\%$	
..... <i>Salix pulchra</i> / <i>Carex aquatilis</i> - <i>Eriophorum angustifolium</i> / <i>Sphagnum</i>	
10b. Vegetation not as above.....	11
11a. <i>Saxifraga hirculus</i> cover $\geq 3\%$	
..... <i>Salix pulchra</i> / <i>Carex aquatilis</i> - <i>Eriophorum angustifolium</i> - <i>Saxifraga hirculus</i>	
11b. Vegetation not as above.....	<i>Salix pulchra</i> / <i>Carex aquatilis</i> - <i>Eriophorum angustifolium</i>
9b. Vegetation not as above.....	12
12a. <i>Carex aquatilis</i> cover $\geq 5\%$	13
13a. <i>Comarum palustre</i> cover $\geq 3\%$	<i>Salix pulchra</i> / <i>Carex aquatilis</i> - <i>Comarum palustre</i>
13b. Vegetation not as above.....	<i>Salix pulchra</i> / <i>Carex aquatilis</i>
12b. Vegetation not as above.....	14
14a. <i>Eriophorum angustifolium</i> $\geq 5\%$	<i>Salix pulchra</i> / <i>Eriophorum angustifolium</i>
14b. Vegetation not as above.....	Undefined Association
6b. Vegetation not as above.....	Undefined Association
5b. Vegetation not as above.....	Undefined Alliance
4b. Sites located in minerotrophic wetlands, i.e., hydrology controlled predominantly by groundwater or surface water from overbank flooding.....	15
15a. Combined cover of low shrubs $\geq 25\%$. Arctic Minerotrophic Wet Low Shrublands Alliance Proposed (A4367p).	16
16a. <i>Salix richardsonii</i> is the dominant low shrub.....	17
17a. Combined forb cover is greater than combined graminoid cover.....	18
18a. <i>Equisetum variegatum</i> cover $\geq 5\%$	<i>Salix richardsonii</i> / <i>Equisetum variegatum</i>

18b. Vegetation not as above.....	Undefined Association
17b. Vegetation not as above.....	19
19a. Combined cover of <i>Carex aquatilis</i> and <i>Eriophorum angustifolium</i> ≥10%.....	
..... <i>Salix richardsonii/Carex aquatilis-Eriophorum angustifolium</i>	
19b. Vegetation not as above.....	Undefined Association
16b. Vegetation not as above.....	Undefined Association
15b. Vegetation not as above.....	Undefined Alliance
3b. Vegetation not as above.....	20
20a. Site permanently flooded with approximately ≥15 cm of water. Sites typically located in small ponds, oxbow lakes, beaded streams, or along lake margins. Not considered further here.....	
..... North American Arctic Freshwater Marsh Group (G370)	
20b. Site not as above. Soils merely wet, or if flooded, then flooded with approximately <15 cm of water, or flooded only early in the freeze-free period. Not considered further here. . North American Arctic Wet Meadow Group (G617)	
1b. The site located in the Arctic and is not influenced by salt water. The soils are moist to dry for most or all of the freeze-free period. The vegetation is dominated by mesomorphic or xeromorphic plants with at least 10% cover of vascular plants, or the site is partially vegetated or barren with <10% cover of vascular plants. Arctic Tundra & Barrens Division (4.B.2.Xa) ..	
.....	21
21a. Site is barren (<10% vascular plant cover) or site is vegetated and located on a river bar, lower floodplain position, or sand dune. M175: Arctic Scree, Rock & Cliff Barrens Macrogroup (M175).....	22
22a. The site is located on a river bar, lower floodplain position, or inland dune. Arctic Gravel Floodplain Vegetation Group (G616).	23
23a. Site is barren (<10% vascular plant cover) or partially vegetated (i.e., vascular plant cover 10–30%) and <i>Salix alaxensis</i> is always present typically at low abundance (<25%). <i>Chamerion latifolium</i> - <i>Salix alaxensis</i> Arctic Floodplain Alliance (A4362).....	24
24a. <i>Salix alaxensis</i> is most the common shrub.	25
25a. <i>Chamerion latifolium</i> cover ≥3%..... <i>Salix alaxensis/Chamerion latifolium</i>	
25b. Vegetation not as above.....	Undefined Association
24b. Vegetation not as above.....	Undefined Association
23b. Vegetation not as above.....	26
26a. <i>Salix alaxensis</i> , <i>S. niphoclada</i> , or <i>S. hastata</i> are the dominant or codominant shrubs.....	27
27a. Sites located predominantly on sand dunes, occassionally on river bars. <i>Salix alaxensis</i> - <i>Salix niphoclada</i> River Bar & Dune Alliance Proposed (A4365p).....	28
28a. <i>Salix niphoclada</i> is the dominant willow, and <i>S. alaxensis</i> cover is ≥5%.....	29
29a. <i>Arctostaphylos rubra</i> cover ≥10%..... <i>Salix niphoclada-Salix alaxensis/Arctostaphylos rubra</i>	
29b. Vegetation not as above.....	Undefined Association
28b. Vegetation not as above.....	30
30a. <i>Salix alaxensis</i> is the dominant willow and <i>Arctostaphylos rubra</i> cover is ≥10%..... <i>Salix alaxensis/Arctostaphylos rubra</i>	
30b. Vegetation not as above.....	31
31a. <i>Salix alaxensis</i> is the dominant willow. <i>Tanacetum bipinnatum</i> ssp. <i>bipinnatum</i> cover is ≥3% and site located on dunes, or <i>T. bipinnatum</i> ssp. <i>bipinnatum</i> cover ≥5% and site located on a river bar.	
..... <i>Salix alaxensis/Tanacetum bipinnatum</i> ssp. <i>bipinnatum</i>	
31b. Vegetation not as above.....	32

32a. <i>Salix alaxensis</i> is the dominant willow and <i>Artemisia campestris</i> ssp. <i>borealis</i> var. <i>borealis</i> cover $\geq 3\%$	<i>Salix alaxensis/Artemisia campestris</i> ssp. <i>borealis</i> var. <i>borealis</i>
..... 32b. Vegetation not as above.	Undefined Association
27b. Sites located on river bars or lower floodplain positions. <i>Salix alaxensis</i> River Bar Alliance Proposed (A4363p).	33
..... 33a. <i>Salix hastata</i> is the dominant willow, and <i>S. alaxensis</i> cover is $\geq 5\%$.	34
..... 34a. <i>Equisetum variegatum</i> cover $\geq 5\%$.	
..... <i>Salix hastata-Salix alaxensis/Equisetum variegatum</i>	
..... 34b. Vegetation not as above.	Undefined Association
33b. Vegetation not as above.	35
..... 35a. <i>Salix alaxensis</i> is the dominant willow.	36
..... 36a. <i>Dryas integrifolia</i> cover is $\geq 10\%$.	<i>Salix alaxensis/Dryas integrifolia</i>
..... 36b. Vegetation not as above.	37
..... 37a. <i>Hedysarum boreale</i> ssp. <i>mackenziei</i> cover $\geq 10\%$.	
..... <i>Salix alaxensis/Hedysarum boreale</i> ssp. <i>mackenziei</i>	
..... 37b. Vegetation not as above.	38
..... 38a. <i>Equisetum arvense</i> cover $\geq 10\%$.	<i>Salix alaxensis/Equisetum arvense</i>
..... 38b. Vegetation not as above.	39
..... 39a. <i>Eurybia sibirica</i> cover $\geq 5\%$.	<i>Salix alaxensis/Eurybia sibirica</i>
..... 39b. Vegetation not as above.	Undefined Association
..... 35b. Vegetation not as above.	Undefined Association
26b. Vegetation not as above.	40
..... 40a. <i>Salix glauca</i> is the dominant or co-dominant shrub. <i>Salix glauca</i> River Bar & Dune Alliance (A4364p).	41
..... 41a. <i>Arctostaphylos rubra</i> cover $\geq 5\%$.	<i>Salix glauca/Arctostaphylos rubra</i>
..... 41b. Vegetation not as above.	42
..... 42a. <i>Koeleria asiatica</i> cover $\geq 3\%$.	<i>Salix glauca/Koeleria asiatica</i>
..... 42b. Vegetation not as above.	Undefined Association
..... 40b. Vegetation not as above.	Undefined Alliance
22b. Site not as above.	43
..... 43a. Site with $\geq 25\%$ lichen cover. Not considered further here.	Arctic Lichen Barrens Group (G868)
..... 43b. Site not as above. Not considered further here.	Arctic Scree, Rock & Cliff Barrens Group (G869)
21b. Site not as above.	44
..... 44a. Site is vegetated ($\geq 10\%$ vascular plant cover) and located on a landform other than a riverbar, lower floodplain position, or inland dune. Arctic Dry-Moist Tundra Macrogroup (M173).	45
..... 45a. Cover of whole tussocks at least 15% and the cover of <i>Eriophorum vaginatum</i> and/or <i>Carex bigelowii</i> $\geq 10\%$. Arctic Herbaceous Tussock Tundra Group Proposed (G899p).	46
..... 46a. Combined cover of low (0.2-1.5 m) and tall (≥ 1.5 m) shrubs $\geq 25\%$ or combined cover of low, tall, and true dwarf shrubs $\geq 35\%$.	47
..... 47a. Soil pH is acidic (<5.7) to circumacidic (5.7-6.5). The shrub component of the vegetation is dominated by alder (<i>Alnus</i>), <i>Betula nana</i> , or ericaceous shrubs. Arctic Acidic Shrub Tussock Tundra Alliance Proposed (A4344p).	48
..... 48a. <i>Alnus viridis</i> ssp. <i>fruticosa</i> cover $\geq 10\%$ and <i>Betula nana</i> cover $\geq 5\%$.	
..... <i>Alnus viridis</i> ssp. <i>fruticosa/Betula nana/Eriophorum vaginatum</i>	
..... 48b. Vegetation not as above.	49

49a. <i>Betula nana</i> cover ≥10%	50
50a. <i>Ledum palustre</i> ssp. <i>decumbens</i> cover ≥10%, or combined cover of <i>L. palustre</i> ssp. <i>decumbens</i> and <i>Vaccinium vitis-idaea</i> ≥15%.	<i>Betula nana-Ledum palustre</i> ssp. <i>decumbens</i> / <i>Eriophorum vaginatum</i>
50b. Vegetation not as above.....	Undefined Association
49b. Vegetation not as above.....	51
51a. <i>Ledum palustre</i> ssp. <i>decumbens</i> cover ≥10% or combined cover of <i>L. palustre</i> ssp. <i>decumbens</i> and <i>Vaccinium vitis-idaea</i> ≥15%.	<i>Ledum palustre</i> ssp. <i>decumbens</i> / <i>Eriophorum vaginatum</i>
51b. Vegetation not as above.....	Undefined Association
47b. Soil pH is circumalkaline (6.6–7.2) to alkaline (>7.2). The shrub component of the vegetation is dominated by <i>Salix pulchra</i> or codominated by <i>S. pulchra</i> and <i>Betula nana</i> . Arctic Nonacidic Shrub Tussock Tundra Alliance Proposed (A4345p).....	52
52a. <i>Betula nana</i> cover ≥10%.....	53
53a. <i>Salix pulchra</i> cover ≥10%.....	<i>Betula nana-Salix pulchra</i> / <i>Eriophorum vaginatum</i>
53b. Vegetation not as above.....	Undefined Association
52b. Vegetation not as above.....	54
54a. <i>Salix pulchra</i> cover ≥10%.....	<i>Salix pulchra</i> / <i>Eriophorum vaginatum</i>
54b. Vegetation not as above.....	Undefined Association
46b. Vegetation not as above.....	55
55a. Soil pH is acidic (<5.7) to circumacidic (5.7–6.5). The shrub component of the vegetation is dominated by ericaceous shrubs. Arctic Acidic Tussock Tundra Alliance Proposed (A4346p).....	56
56a. <i>Vaccinium uliginosum</i> cover ≥5% or combined cover of <i>V. uliginosum</i> and other ericaceous shrubs ≥10% and <i>Sphagnum</i> cover ≥10%.....	<i>Eriophorum vaginatum</i> / <i>Vaccinium uliginosum</i> / <i>Sphagnum</i>
56b. Vegetation not as above.....	57
57a. Combined cover of <i>Ledum palustre</i> ssp. <i>decumbens</i> and <i>Vaccinium vitis-idaea</i> ≥10%, or combined cover of <i>L. palustre</i> ssp. <i>decumbens</i> and <i>V. vitis-idaea</i> and other ericaceous shrubs ≥15%.....	<i>Eriophorum vaginatum</i> / <i>Ledum palustre</i> ssp. <i>decumbens</i> - <i>Vaccinium vitis-idaea</i>
57b. Vegetation not as above.....	Undefined Association
55b. Soil pH is circumalkaline (6.6–7.2) to alkaline (>7.2). The shrub component of the vegetation is dominated by the prostrate shrubs <i>Dryas integrifolia</i> , <i>Salix reticulata</i> , and/or <i>Arctostaphylos rubra</i> . Arctic Nonacidic Tussock Tundra Alliance Proposed (A4347p).....	58
58a. <i>Dryas integrifolia</i> cover ≥5%.....	<i>Eriophorum vaginatum</i> / <i>Dryas integrifolia</i>
58b. Vegetation not as above.....	Undefined Association
45b. Vegetation not as above.....	59
59a. Combined cover of low (0.2–1.5 m) and tall (≥1.5 m) shrubs ≥25% or combined cover of low, tall, and true dwarf shrubs ≥35%	60
60a. Combined cover of low (0.2–1.5 m) and tall (≥1.5 m) shrubs ≥25%. Arctic Low Shrub Tundra (G897).	61
61a. <i>Betula nana</i> is the dominant shrub, or <i>B. nana</i> is codominant with low willows (<i>Salix</i>). Arctic Dwarf Birch Low Shrub Tundra Alliance (A4339).	62
62a. <i>Salix glauca</i> is codominant with <i>B. nana</i>	63
63a. Cover of <i>Vaccinium vitis-idaea</i> ≥10% or combined cover of ericaceous shrubs ≥15% and <i>Vaccinium vitis-idaea</i> ≥3%.....	64
64a. <i>Carex bigelowii</i> cover ≥5%.	<i>Betula nana-Salix glauca</i> / <i>Vaccinium vitis-idaea</i> / <i>Carex bigelowii</i>

64b. Vegetation not as above.....	65
65a. <i>Saussurea angustifolia</i> cover $\geq 3\%$	
..... <i>Betula nana-Salix glauca/Vaccinium vitis-idaea/Saussurea angustifolia</i>	
65b. Vegetation not as above.....	Undefined Association
63b. Vegetation not as above.....	Undefined Association
62b. Vegetation not as above.....	66
66a. <i>Salix pulchra</i> is codominant with <i>B. nana</i>	67
67a. <i>Petasites frigidus</i> cover $\geq 5\%$	<i>Betula nana-Salix pulchra/Petasites frigidus</i>
67b. Vegetation not as above.....	Undefined Association
66b. Vegetation not as above.	Undefined Association
61b. Vegetation not as above.....	68
68a. Low willows are the dominant shrub.....	69
69a. Soil pH is acidic (<5.7) to circumacidic (5.7–6.5). Arctic Acidic Low Willow Tundra Alliance (A4337).....	
..... 70	
70a. <i>Salix pulchra</i> is the dominant willow.....	71
71a. Combined cover of ericaceous shrubs $\geq 15\%$	72
72a. <i>Vaccinium uliginosum</i> cover $\geq 5\%$	<i>Salix pulchra-Vaccinium uliginosum</i>
72b. Vegetation not as above.....	Undefined Association
71b. Vegetation not as above.....	73
73a. Combined forb cover is greater than combined graminoid cover.....	74
74a. <i>Equisetum arvense</i> cover $\geq 5\%$	<i>Salix pulchra/Equisetum arvense</i>
74b. Vegetation not as above.....	75
75a. <i>Petasites frigidus</i> cover $\geq 5\%$	<i>Salix pulchra/Petasites frigidus</i>
75b. Vegetation not as above.....	Undefined Association
73b. Vegetation not as above.....	76
76a. <i>Carex bigelowii</i> cover $\geq 5\%$	<i>Salix pulchra/Carex bigelowii</i>
76b. Vegetation not as above.....	77
77a. <i>Hylocomium splendens</i> cover $\geq 25\%$	<i>Salix pulchra/Hylocomium splendens</i>
77b. Vegetation not as above.....	Undefined Association
70b. Vegetation not as above.....	Undefined Association
69b. Soil pH is circumalkaline (6.6–7.2) to alkaline (>7.2). Arctic Nonacidic Low Willow Tundra Alliance (A4338).....	78
78a. <i>Salix glauca</i> is the dominant willow.....	79
79a. Cover of <i>Dryas integrifolia</i> $\geq 10\%$ or <i>D. integrifolia</i> cover $\geq 3\%$ and combined cover of dwarf willows and <i>D. integrifolia</i> $\geq 15\%$	80
80a. <i>Carex bigelowii</i> cover $\geq 5\%$	<i>Salix glauca/Dryas integrifolia/Carex bigelowii</i>
80b. Vegetation not as above.....	81
81a. <i>Rhytidium rugosum</i> cover $\geq 3\%$	<i>Salix glauca/Dryas integrifolia/Rhytidium rugosum</i>
81b. Vegetation not as above.....	Undefined Association
79b. Vegetation not as above.....	82
82a. Combined forb cover is greater than combined graminoid cover.....	83
83a. <i>Lupinus arcticus</i> cover $\geq 5\%$	<i>Salix glauca/Lupinus arcticus</i>
83b. Vegetation not as above.....	Undefined Association

82b. Vegetation not as above.....	84
84a. <i>Arctagrostis latifolia</i> cover ≥5%.....	<i>Salix glauca/Arctagrostis latifolia</i>
84b. Vegetation not as above.....	Undefined Association
78b. Vegetation not as above.....	85
85a. <i>Salix richardsonii</i> is the dominant willow	86
86a. <i>Arctostaphylos rubra</i> cover ≥5%.....	<i>Salix richardsonii/Arctostaphylos rubra</i>
86b. Vegetation not as above	87
87a. Combined forb cover is greater than combined graminoid cover.....	88
88a. <i>Equisetum arvense</i> cover ≥5%	89
89a. <i>Festuca altaica</i> cover ≥3%.....	<i>Salix richardsonii/Equisetum arvense-Festuca altaica</i>
89b. Vegetation not as above.....	90
90a. <i>Petasites frigidus</i> cover ≥3%.....	
.....	<i>Salix richardsonii/Equisetum arvense-Petasites frigidus</i>
90b. Vegetation not as above.....	<i>Salix richardsonii/Equisetum arvense</i>
88b. Vegetation not as above.....	Undefined Association
87b. Vegetation not as above.....	Undefined Association
85b. Vegetation not as above.....	Undefined Association
68b. Vegetation not as above.....	Undefined Alliance
60b. Vegetation not as above. Not considered further here.....	Arctic Dwarf Shrub Tundra Group (G896)
59b. Vegetation not as above. Not considered further here.....	
.....	Arctic Herbaceous Nontussock Tundra Group Proposed (G898p)
44b. Vegetation and/or site not as above.....	Undefined Macrogroup